



# STIMUTER



## Roots in balance for strawberries

(Growing on gutters, Fruit growing greenhouse and/or tunnel fruit growing)

**STIMUTER** can be used when abiotic stress can occur due to external factors and a less vital soil/substrate and weak resilience of plants and crops. The product supports the natural regenerative capacity, has a nourishing, nurturing and plant-strengthening effect in case of external influences, in the soil and at the roots.

**STIMUTER** is absorbed by the soil, leaf and roots of the plant. Thanks to the available bio-stimulant and crop care components, plants will develop increased resilience and vitality.



After planting  
10L/ha Stimuter  
lq.

After 2  
weeks 10L/ha  
Stimuter lq.

After 2  
weeks 10L/ha  
Stimuter lq.

After 2  
weeks 10L/ha  
Stimuter lq.

After 2  
weeks 10L/ha  
Stimuter lq.

**For optimal performance of Stimuter, leaf application is ideal.**

**Stimuter can be dripped and/or used as a foliar spraying.**

**Administration of Stimuter liquid preferably in the early morning or evening.**

**Can also be used before irrigation.**

- Carry out a test spray before use.
- Do not use the product with bright sunlight.
- Pireco cannot be held liable for any damage resulting from improper use.
- Always consult your cultivation advisor for suitable advice and combination applications.